

MSHSM – Minimal Standard Heterotic String Models

Alon E. Faraggi^{1*}

¹ Department of Mathematical Sciences, University of Liverpool, Liverpool L69 7ZL, UK

Key words String phenomenology, free fermionic models

An overview of old and new results in studies of the quasi-realistic free fermionic models is presented, which include the recent discovery of exophobic string vacua and reproduction of the Higgs–matter splitting mechanism in a corresponding orbifold construction.

Copyright line will be provided by the publisher

Contemporary terrestrial experiments, and extra-terrestrial observatories, indicate that the Standard $SU(3)_C \times SU(2)_L \times U(1)_Y$ Model of the gauge interactions, augmented with three families of quarks and leptons, correctly accounts for the extracted data. The remaining piece, the scalar Higgs field and the mechanism of electroweak symmetry breaking, awaits the data and final judgement of the LHC experiment. Post 1998 data from astrophysical and reactor neutrino experiments also indicate that neutrinos are massive, which suggests augmentation of each Standard Model generation with a Standard Model singlet.

Perhaps its most tantalising property, is the fact that the Standard Model matter and gauge sectors can be embedded into Grand Unified Theories. Most appealing in this context is the embedding into $SO(10)$, that accomodates each matter generation into a single spinorial 16 representation. The gauge charges of the matter states are experimental observables and the embedding into $SO(10)$ reduces these number of observables from $54 = 3 \times 3 \times 6$, to just one, namely the number of generations. This could of course be a cruel accident, but it is the best clue we have for the fundamental origins of the Standard Model.

Additional evidence for the validity of grand unification exists in the logarithmic running of the Standard Model parameters, which is compatible with the data in the gauge and heavy matter sectors. The Higgs sector receives quadratic corrections from the cutoff scale, and restoration of its logarithmic evolution requires the introduction of a new symmetry – *e.g.* supersymmetry. Additionally, proton longevity and neutrino masses suggest that the unification scale is vastly separated from the electroweak scale. Indeed, proton stability is an accidental feature of the Standard Model at the renormalizable level. Non-renormalizable operators, suppressed by an effective cutoff scale, which must exist because the Standard Model is incomplete, will, in general, induce fast proton decay. The proton lifetime is therefore another fact, set in our bones, that hints at a large unification scale.

Gran unification, however, cannot be the end of the story. The flavour data is not explained in this context. The choice of gauge and matter sectors is also ad hoc. Most importantly, gravity, is not included in this framework. Furthermore, non-conformal quantum field theories and general relativity are incompatible at a fundamental level. String theory is a unique extension of the Standard Model in the sense that while providing a consistent theory of quantum gravity, it also gives rise to the gauge and matter structures that appear in the Standard Model. It provides a unifying framework for all the basic particle constituents that appear in nature, and enables the development of a phenomenological approach to quantum gravity. Furthermore, string theory is predictive. It predicts that the rank of the gauge interactions is extended, and should be, for example, below 22 in the heterotic string.

Given that there are five different theories in ten dimensions, which are all believed to be connected by various duality transformations, a natural question is which of these theories should be used as a starting point for phenomenology. The Standard Model data motivates the hypothesis that the perturbative string

* Corresponding author E-mail: alon.faraggi@liv.ac.uk, Phone: +00 44 0151 7943774, Fax: +00 44 0151 7943784

vacuum should accommodate the canonical $SO(10)$ embedding of its spectrum. Additionally, a perturbative string vacuum should contain three chiral generations and a pair of Higgs doublets with potentially viable Yukawa couplings. Among the five perturbative string theories the only one that is compatible with these requirements is the $E_8 \times E_8$ heterotic-string, as it is the only one that produces spinorial representations in the perturbative massless spectrum. It may well be that exploring other properties of the fundamental vacuum will necessitate utilising a different perturbative limit of the underlying non-perturbative theory. The ultimate question in this regard is how are the different limits connected, and how can we be guaranteed that we are exploring different limits of a single nonperturbative vacuum.

The quasi-realistic free fermionic heterotic-string models were constructed in the late eighties and early nineties [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. They provide a concrete framework to study many of the issues that pertain to the phenomenology of the Standard Model and grand unification. A few highlights of studies in the free fermionic standard-like models are listed below:

- Top quark mass $\sim 175\text{--}180\text{GeV}$ [6]
- Generation mass hierarchy [14]
- CKM mixing [15, 16]
- Stringy seesaw mechanism [17, 18, 19]
- Gauge coupling unification [20, 21]
- Proton stability [22]
- Squark degeneracy [23, 24]
- Minimal Standard Heterotic String Model (MSHSM) [8]
- Moduli fixing [25]
- Classification [26, 27] & spinor-vector duality [28, 29]

The fermionic formulation was developed in the mid-eighties [13, 30]. The equivalence of bosons and fermions in two dimensions entails that a model constructed using the fermionic approach corresponds to a model constructed using the bosonic approach in which the target-space is compactified on a six dimensional internal manifold. The free fermionic formalism corresponds to using a free bosonic formalism in which the radii of the internal dimensions are fixed at a special point in the compact space. Deformation from the special point in the moduli space are parametrized in terms of world-sheet Thirring interactions among the world-sheet fermions [31]. The simplicity of the free fermionic formalism entails that the string consistency constraints are solved in terms of the world-sheet free fermion transformation properties on the string world-sheet, which are encoded in sets of basis vectors and one-loop GSO projection coefficients among the basis vectors. The formalism to extract the physical spectrum and superpotential interaction terms are also straightforward. The simplest free fermionic constructions correspond to a $Z_2 \times Z_2$ orbifold of a six dimensional toroidal manifold, augmented with discrete Wilson lines that are needed to break the $SO(10)$ GUT symmetry.

Perhaps, the most tantalising achievement is the successful calculation of the top quark mass [6], which was obtained several years prior to the experimental discovery. This calculation demonstrates how string theory enables the calculation of the fermion-scalar Yukawa couplings in terms of the unified gauge coupling. Furthermore, the string models offer an explanation for the hierarchical mass splitting between the top and bottom quarks. The top quark Yukawa coupling is obtained at the cubic level of the superpotential and is of order one, whereas the Yukawa couplings of the lighter quarks and leptons are obtained from nonrenormalizable operators that are suppressed relative to the leading cubic level term. Thus, only the top quark mass is characterised by the electroweak scale and the masses of the lighter quarks and leptons are

naturally suppressed compared to it. As the heavy generation Yukawa couplings are obtained at low orders in the superpotential, the calculation of these Yukawa couplings is robust and is common to a large class of models. Further analysis of fermion masses shows that quasi-realistic fermion mass textures arise for reasonable choices of supersymmetric flat directions [15]. Issues like left-handed neutrino masses [18], gauge coupling unification [20, 21], proton stability [22] and squark degeneracy [23, 24] were studied in concrete quasi-realistic free fermionic string models and for detailed solutions of the supersymmetric flat direction constraints. It was also demonstrated in ref. [8] that the free fermionic heterotic string vacua give rise to models that produce in the observable charged sector below the string unification scale solely the matter spectrum of the minimal supersymmetric standard model. Such models are dubbed Minimal Standard Heterotic String Models (MSHSM). The free fermionic models also provide important clues to the problem of moduli fixing in string theory [25]. They highlight the fact that string theory may utilize geometrical structures that do not have a classical correspondence. Primarily, they allow boundary conditions that distinguish between the left- and right-moving coordinates of the six dimensional compactified space. Such boundary conditions necessarily lead to the projection of the moduli fields associated with the extra internal coordinates. The free fermionic models have also been instrumental in recent years to unravel a new duality symmetry under the exchange of spinor and vector representations of the GUT group [28, 29]. The free fermionic formalism is also adaptable to a computerised classification, of $Z_2 \times Z_2$ orbifolds with symmetric shifts [26, 27]. A generic property of string vacua is the existence of fractionally charged states, due to the breaking of the $SO(10)$ GUT symmetry by discrete Wilson lines. The lowest state among those is stable due to electric charge conservation. Experimental constraints severely constrain the mass and abundance of such particles. Using the free fermionic classification method we recently constructed examples of Pati-Salam string models that are completely free of massless exotics [32].

The quasi-realistic free fermion models were constructed in the late eighties and early nineties, and continue to be represent viable string vacua that merit deeper exploration into their properties. In particular, their relation to $Z_2 \times Z_2$ orbifolds is of vital importance. One should note that the different formalisms of string compactifications are related and provide complementary insight to the properties of the underlying vacua. So, for example, while the free fermionic approach provides easy access to the massless spectra and potential, and is adaptable to a computerised classification, orbifold representation may be more amenable to study moduli dynamics and vacuum selection. In this vein, it is vital to establish a detailed dictionary between the two languages. While each language may be well understood on its own, writing a detailed dictionary between the two is often non-trivial.

In this talk I highlight how this correspondence can be established for two key properties of the models, and is indeed found to be highly non-trivial. The first is with respect to the $Z_2 \times Z_2$ orbifold action on the $SO(12)$ lattice at the free fermionic point. It is well known that a $Z_2 \times Z_2$ orbifold of a generic six dimensional torus produces 48 fixed points, whereas $Z_2 \times Z_2$ of the $SO(12)$ lattice produces 24 fixed points. The 48 points can be reduced to 24 by adding a freely acting shift, under which pairs of fixed points are identified. Now one can add different freely acting shifts to this effect, and the question is which one reproduces the free fermionic model. Namely, it is not sufficient to examine the massless spectrum for this purpose, but one must also establish the correspondence of the massive spectrum. String theory affords even more interesting possibilities because, in addition to the ordinary Kaluza-Klein momentum modes, it also gives rise to space-like winding modes. Therefore, one can add freely acting shifts that shift the momenta, winding, as well as asymmetric shifts that act on both the momenta and winding and are therefore intrinsically stringy! It turns out that the shifts that reproduce the partition function at the free fermionic point are uniquely of the third kind.

The second property, related to the free fermion-orbifold correspondence, illustrates the utility of the free fermion method in accessing classes of string vacua that are obscure from the point of view of the orbifold construction. In the orbifold models one typically starts with the $E_8 \times E_8$ heterotic-string compactified to four dimensions and subsequently break one E_8 gauge factor by the orbifold twistings and discrete Wilson lines. In contrast the quasi-realistic free fermionic models start with $SO(16) \times SO(16)$, where the reduction from $E_8 \times E_8$ to $SO(16) \times SO(16)$ is realised in terms of a Generalised GSO (GGSO) phase in the partition function. One can then show that the partition function at the free fermionic point

is generated by starting with the partition function of the $E_8 \times E_8$ and acting on it with a freely acting $Z_2 \times Z_2$ twist given by $Z_2^a \times Z_2^b := (-1)^{F_{\xi^1} \delta_1} \times (-1)^{F_{\xi^2} \delta_1}$ with $\delta_1 X^9 = X_9 + \pi R_9$, where $F_{\xi^{1,2}}$ are fermion numbers acting on the first and second E_8 factors, and δ_9 is a mod 2 shift in one internal direction [33]. The action of the $Z_2 \times Z_2$ freely acting twist reduces the $E_8 \times E_8$ symmetry to $SO(16) \times SO(16)$. The next step in the construction employs a Z_2^c twist that acts on the internal coordinates and produces 16 fixed points. The gauge symmetry is broken to $E_7 \times SU(2)$, and $SO(12) \times SO(4)$, in each case, respectively. The matter and Higgs states in the first case are in the 56 representation of E_7 , which breaks as $56 = (32, 1) + (12, 2)$ under $SO(12) \times SU(2)$, where the 32 spinorial, and the 12 vectorial, representations of $SO(12)$ contain the matter and Higgs representations of the Standard Model, respectively. Hence, the freely acting $Z_2^a \times Z_2^b$ twists induce a string Higgs–matter splitting mechanism, similar to the string doublet–triplet splitting mechanism [22]. However, it turns out that simply adding the Z_2^c twist, *i.e.* taking $[Z_+/(Z_2^a \times Z_2^b)]/Z_2^c$, where Z_+ is the $E_8 \times E_8$ partition function, projects the twisted massless spinorial representations and keeps the vectorials. To restore the matter representations one needs to analyse the full $Z_+/(Z_2^a \times Z_2^b \times Z_2^c)$ partition and take into account the seven independent modular orbits (seven discrete torsions) [35]. A non–trivial proposition indeed! Now, one should note that this is merely one example. The free fermionic classification accesses 2^5 vacua at this level, and by breaking the hidden $SO(16)$ factor to $SO(8) \times SO(8)$ this is increased to 2^9 . We therefore note that the free fermionic approach provides a superior tool to easily access large classes of string vacua. The bosonic formalism provides a more intuitive approach to the string moduli and hopefully to the dynamics that fix the string vacuum.

Acknowledgements This work is supported in part by the STFC under grant ST/G00062X/1 and the EU under contract MRTN-CT-2006-035863.

References

- [1] I. Antoniadis, J. Ellis, J. Hagelin and D.V. Nanopoulos *Phys. Lett. B* **231**, 65 (1989).
- [2] A.E. Faraggi, D.V. Nanopoulos and K. Yuan, *Nucl. Phys. B* **335**, 347 (1990).
- [3] I. Antoniadis, G.K. Leontaris and J. Rizos, *Phys. Lett. B* **245**, 161 (1990).
- [4] A.E. Faraggi and D.V. Nanopoulos, *Phys. Rev. D* **48**, 3288 (1993).
- [5] A.E. Faraggi, *Phys. Lett. B* **278**, 131 (1992); *Nucl. Phys. B* **403**, 101 (1993).
- [6] A.E. Faraggi, *Phys. Lett. B* **274**, 47 (1992); *Phys. Lett. B* **339**, 223 (1994).
- [7] A.E. Faraggi, *Nucl. Phys. B* **387**, 239 (1992).
- [8] G.B. Cleaver, A.E. Faraggi and D.V. Nanopoulos, *Phys. Lett. B* **455**, 135 (1999).
- [9] G.B. Cleaver, A.E. Faraggi and C. Savage, *Phys. Rev. D* **63**, 066001 (2001).
- [10] G. Cleaver, A.E. Faraggi and S. Nooij, *Nucl. Phys. B* **672**, 64 (2003).
- [11] A.E. Faraggi, E. Manno and C. Timirgaziu, *Eur. Phys. Jour. C* **50**, 701 (2007).
- [12] G. Cleaver, A.E. Faraggi, E. Manno and C. Timirgaziu, *Phys. Rev. D* **78**, 046009 (2008).
- [13] I. Antoniadis, C. P. Bachas, and C. Kounnas, *Nucl. Phys. B* **289**, 87 (1987).
- [14] A.E. Faraggi, *Nucl. Phys. B* **407**, 57 (1993).
- [15] I. Antoniadis, J. Rizos and K. Tamvakis, *Phys. Lett. B* **278**, 257 (1992).
- [16] A.E. Faraggi and E. Halyo, *Phys. Lett. B* **307**, 305 (1993); *Nucl. Phys. B* **416**, 63 (1994).
- [17] I. Antoniadis, J. Rizos and K. Tamvakis, *Phys. Lett. B* **279**, 281 (1992).
- [18] A.E. Faraggi and E. Halyo, *Phys. Lett. B* **307**, 311 (1993).
- [19] C. Coriano and A.E. Faraggi, *Phys. Lett. B* **581**, 99 (2004).
- [20] A.E. Faraggi, *Phys. Lett. B* **302**, 202 (1992).
- [21] K.R. Dienes and A.E. Faraggi, *Phys. Rev. Lett.* **75**, 2646 (1995); *Nucl. Phys. B* **457**, 409 (1995).
- [22] A.E. Faraggi, *Nucl. Phys. B* **428**, 111 (1994); *Phys. Lett. B* **499**, 147 (2001); *Phys. Lett. B* **520**, 337 (2001).
- [23] A.E. Faraggi and J.C. Pati, *Nucl. Phys. B* **526**, 21 (1998).
- [24] A.E. Faraggi and O. Vives, *Nucl. Phys. B* **641**, 93 (2002).
- [25] A.E. Faraggi, *Nucl. Phys. B* **728**, 83 (2005).
- [26] A.E. Faraggi, C. Kounnas, S.E.M. Nooij and J. Rizos, *Nucl. Phys. B* **695**, 41 (2004).
- [27] A.E. Faraggi, C. Kounnas and J. Rizos, *Phys. Lett. B* **648**, 84 (2007);
- [28] A.E. Faraggi, C. Kounnas and J. Rizos, *Nucl. Phys. B* **774**, 208 (2007); *Nucl. Phys. B* **799**, 19 (2008).
- [29] T. Catelin-Jullien, A.E. Faraggi, C. Kounnas and J. Rizos, *Nucl. Phys. B* **812**, 103 (2009).
- [30] H. Kawai, D.C. Lewellen, and S.H.-H. Tye, *Nucl. Phys. B* **288**, 1 (1987).
- [31] S. Elitzur, E. Gross, E. Rabinovici and N. Seiberg, *Nucl. Phys. B* **283**, 413 (1987).
- [32] B. Assel, K. Christodoulides, A.E. Faraggi, C. Kounnas and John Rizos, *Phys. Lett. B* **683**, 306 (2010).
- [33] A.E. Faraggi and M. Tsulaia, *Phys. Lett. B* **683**, 314 (2010).
- [34] A.E. Faraggi, *Eur. Phys. Jour. C* **49**, 803 (2007).
- [35] C. Angelantonj, A.E. Faraggi and M. Tsulaia, paper in preparation.